

Sentri[™] end tidal CO₂ monitoring range

Masks and nasal cannula



Oxygen and Aerosol Therapy • Variable Oxygen Concentration (Low Flow)





Quality, innovation and choice



Sentri[™] end tidal CO₂ monitoring range

Capnography is vital during sedation

The increased use of conscious sedation has created a need for devices to monitor respiratory depression. The difference between conscious sedation and general anaesthesia is sometimes very small. It is possible during conscious sedation that intravenous sedatives and narcotics administered to allay apprehension can result in the loss of consciousness and respiratory obstruction.

Mask or nasal cannula the choice is yours

Sentri is available as an adult mask and in three sizes of nasal cannula. Both permit the sampling of exhaled carbon dioxide in non-intubated patients during the administration of supplementary oxygen.



By delivering oxygen through one prong and sampling exhaled gas from the other, the nasal cannula can provide end tidal CO₂ values comparable to those achieved with intubated patients. Nasal cannula may be more appropriate for paediatric patients when high oxygen flows may "dilute" the CO₂ sample and give a low (or no) value. A face mask may be more appropriate when the nares are occluded or obstructed.

200

Sentri™ Intersurgical EcoLite[™]mask kit

The Sentri Intersurgical EcoLite[™] mask has been designed as part of our ongoing focus on sustainable development, which is an integral part of our ethos.

The Intersurgical EcoLite range of masks provide an option with a significantly lower impact on the environment. By utilising the latest technology we have combined two materials: the first forming the clear rigid shell of the mask, this provides a strong lightweight body with good visibility, whilst the second softer material forms the seal, providing a comfortable fit for a wide variety of face shapes.

Incurved nose seal

Conforms to different nose shapes designed to prevent oxygen entering patient's eyes

No metal nose clip MRI compatible

Multi-channel oxygen tube

Oxygen still flows even if tube is kinked

Luer lock port

Secure ETCO₂ monitoring line connection

Elastic can be positioned under or over the ears Below ear position eliminates trauma

Soft face seals


to top of ears

Increased patient comfort

A choice of "under chin" or "on chin" positions

Provides a better fit on a wider range of patient face shapes

Two integral chin seals ensure the mask fits a wider range of patient face shapes





Comfortable for the environment

Various groups have questioned the use of PVC in medical products and it's impact on the environment. Intersurgical's aim is to reduce the environmental impact of its products and processes. The utilisation of new materials and the latest manufacturing technology has resulted in the elimination of PVC from the mask shell resulting in a reduced environmental impact.





www.intersurgical.com/info/eco

Sentri[™] Intersurgical EcoLite[™] mask

Code		Description	Tube length	Box Qty.	
1142018	5 NEW	Sentri Intersurgical EcoLite ^{TM} , adult, mask with CO ₂ monitoring line and tube	2.1m	30	G
114301	5 NEW	Sentri Intersurgical EcoLite TM , adult, mask with \rm{CO}_2 monitoring line, filter and tube	2.1m	30	

Sentri nasal cannula

Code	Description	Tube length	Box Oty.
1144001	Sentri, adult, nasal cannula with curved prongs and tube	2.1m	50
1144002	Sentri, adult, nasal cannula with curved prongs, CO_{2} monitoring line, filter and tube	2.1m	40
1144009	Sentri, infant, nasal cannula with curved prongs and tube	2.1m	50
1144010	Sentri, infant, nasal cannula with curved prongs, $\mathrm{CO}_{_2}$ monitoring line, filter and tube	2.1m	40
1144005	Sentri, paediatric, nasal cannula with curved prongs and tube	2.1m	50
1144006	Sentri, paediatric, nasal cannula with curved prongs, $\mathrm{CO}_{\rm 2}$ monitoring line, filter and tube	2.1m	40

Lower environmental impact product

References

Venkatesh Srinivasa & Bhavani Shankar Kodali. Miner JR, Heegaard W, Plummer D: End Tidal Carbon Dioxide Monitoring of Procedural Sedation SAEM Scientific Assembly, May 2001. Accurate Determination of End-Tidal Carbon Dioxide During Administration of Oxygen by Nasal Cannulae by Edwin A Bowe, MD; Philip G. Boysen, MD; Julie A. Broome, BS; E.F. Klein, Jr., MD J Clin Monit 1989; 5:105-110 The society for pediatric sedation -sedation provider course.



Intersurgical Ltd, Crane House, Molly Millars Lane, Wokingham, Berkshire, RG41 2RZ, UK T: +44 (0)118 9656 300 F: +44 (0)118 9656 356 info@intersurgical.com www.intersurgical.com

UK · Ireland · France · Germany · Spain · Portugal Italy · Benelux · Sweden · Denmark · Lithuania · Russia Czech Republic • Turkey • South Africa • China • Japan Taiwan · Philippines · USA · Canada · Colombia · Australia







IS10.24 • Issue 5 02.17

ΜΙΧ

FSC* C013343

From respons sources

Intersurgical Ltd is certified to ISO 14001, ISO 9001 and ISO 13485